

3. Haynes AB, Deshpande V, Ingkakul T, Vagefi PA, Szymonifka J, Thayer SP, et al. Implications of incidentally discovered, non-functioning pancreatic endocrine tumors: short-term and long-term patient outcomes. *Arch Surg.* 2011;146(5):534-8.
4. Lee LC, Grant CS, Salomao DR, Fletcher JG, Takahashi N, Fidler JL, et al. Small, non-functioning, asymptomatic pancreatic neuroendocrine tumors (PNETs): role for non-operative management. *Surgery.* 2012;152(6):965-74.
5. Toste PA, Kadera BE, Tatishchev SF, Dawson DW, Clerkin BM, Muthusamy R, et al. Nonfunctional Pancreatic Neuroendocrine Tumors <2cm on Preoperative Imaging are Associated with a Low Incidence of Nodal Metastasis and an Excellent Overall Survival. *J Gastrointest Surg.* 2013;8.
6. Gaujoux S, Partelli S, Maire F, D'Onofrio M, Larroque B, Tamburrino D, et al. Observational study of natural history of small sporadic non-functioning pancreatic neuroendocrine tumors. *J Clin Endocrinol Metab.* 2013;20.
7. Kuo EJ, Salem RR. Population-level analysis of pancreatic neuroendocrine tumors 2cm or less in size. *Ann Surg Oncol.* 2013;20(9):2815-21.
8. Bilimoria KY, Talamonti MS, Tomlinson JS, Stewart AK, Winchester DP, Ko CY, et al. Prognostic score predicting survival after resection of pancreatic neuroendocrine tumors: analysis of 3851 patients. *Ann Surg.* 2008;247(3):490-500.
9. Zerbi A, Capitanio V, Boninsegna L, Pasquali C, Rindi G, Delle Fave G, et al. Surgical treatment of pancreatic endocrine tumours in Italy: results of a prospective multicentre study of 262 cases. *Langenbecks Arch Surg.* 2011;396(3):313-21.

---

## Symposium 4 (Video, How I do it?) Efforts to obtain more safety margin during surgery for Klatskin tumor

---

### 1. Conventional resection

Department of Surgery, Seoul National University  
College of Medicine, Korea

**Dae Wook Hwang**

---

#### Bile duct excision

All patients underwent a resection of the extra-hepatic biliary tree from the level of the upper border of the pancreas or about 1cm into intra-pancreatic portion to the confluence of the left and right hepatic ducts. If the tumour extended lower than the intrapancreatic portion of the common bile duct, this was also removed and pancreaticoduodenectomy was considered. At the upper end of the biliary tree, the extent of biliary resection has depended on the extent of hepatic resection required. Routine frozen-section histopathology has also been used to guide the extent of biliary resection. Care is taken to control bile spillage during resection in order to decrease the risk of postoperative tumor seeding. After the distal portion of the bile duct was cut near the edge of the pancreas early during the operation, the stump of the bile duct was meticulously closed to avoid bile spill. Secondly, the proximal, segmental bile ducts were cut in the liver remnant, preferably as the final step in the procedure, after the parenchymal transection has been completely executed, with application of clamp at the tumor-side. Biliary reconstruction in all patients was performed by retrocolic Roux-en-Y hepaticojejunostomy without biliary stents.

#### Hepatic resection

Hepatic resection was carried out to include all segments of liver involved by tumour or affected

by planned loss of vascular supply during surgery. Caudate lobectomy was planned and carried out routinely. Included hepatic segments were S1, S4a and S5-8.

Hepatic transection was accomplished using modified hanging maneuver with a Kelly clamp crushing technique. A small plastic tube was placed retroheptically in the cut plane. The top of the tube was placed between the middle and right hepatic veins. Then, the tube was placed along the border between the left lateral section and Spiegel's caudate lobe (along ductus venosus), and the bottom of the tube was placed at the right side of the umbilical Glissonian pedicle (or the medial side of portal vein of the umbilical portion). During the parenchyma transection, the cut plane will always target to the tube under lifting up the tube by hanging. The left bile duct remained the last step to transect. The area of proximal liver hilum remained untouched in the whole procedure and will be resected en bloc after liver parenchyma transection.

#### Vascular resection

Concomitant portal vein resection and reconstruction have been applied aggressively when macroscopic invasion was suspected at surgery. As some controversial problems remain, concomitant hepatic artery resection and reconstruction have not been applied similarly as portal vein resection. The portal vein will be transected and anastomosed in the early phase of skeletonization of hepatoduodenal ligament, and reconstruction was performed by end-to-end anastomosis or the use of inter-position graft, bovine pericardial patch or PTFE, in specific cases.

#### Lymphadenectomy

Lymphadenectomy has included lymph nodes, lymphatic channels and nerves surrounding the portal vein and hepatic artery in all patients. All the regional lymphatic tissues were dissected and the sampling of para-aortic nodes (No. 16 a2 & b1)

and retropancreatic lymph nodes (No. 13) was taken in most patients. In elderly patients (aged > 75 years) or patients with significant cardiopulmonary comorbidity, lymphadenectomy has been limited to the regional nodes in an attempt to reduce the risk for perioperative mortality.

---

## 2. Extended resection

Center for Liver Cancer, National Cancer Center,  
Korea

**Sang-Jae Park**

---



---

## 3. Limited resection

Department of Surgery, University of Ulsan College  
of Medicine, Korea

**Shin Hwang**

---

Both curative resection and acceptably low perioperative mortality offer the chance of long-term survival in patients with hilar cholangiocarcinoma. The resectability rate for hilar cholangiocarcinoma has increased through concurrent major hepatectomies, but these procedures are technically demanding and often associated with serious complications. Parenchyma-preserving limited hepatectomy (PPLH) has a definite merit on patient safety, but surgical procedure becomes more complex.

PPLH was performed in 15 patients with advanced perihilar cholangiocarcinoma by a single surgeon. Their medical records were analyzed retrospectively.

The types of PPLH included resection of S4+1 (n=10), S5+4a+1(n=2), S5+4+1 (n=2) and isolated S1 (n=1). R0 resection was achieved in 12 patients. Concurrent pylorus-preserving pancreatectomy and extended bile duct resection with excavation of the intrapancreatic bile duct was performed in 3 and 5